



HLI AMERICA

PROTECTING LIFE BEGINS AT HOME

A Program of Human Life International

50 Years of the Pill

Shocking Statistics

- **Prior to 'The Pill' (hormonal contraceptive), there were known to be 5 sexually transmitted diseases (STDs). Today there are more than 30.** Nearly 20 million individual cases of STDs are reported in the U.S. every year, half of which infect 15 – 24 year olds. Over 50 million U.S. men and women are reported to have incurable genital herpes.¹
- **Women who use a hormonal contraceptive for a minimum of four years prior to their first full term pregnancy have a 52% higher risk of developing breast cancer.**²
- **Women who use a hormonal contraceptive for more than five years are four times more likely to develop cervical cancer.**³
- Oral contraceptives are declared Group I Carcinogenic to Humans. A carcinogen is any substance that is an agent directly involved in causing cancer.⁴
- The 'abortifacient action' in hormonal contraceptives is documented by pharmaceutical manufacturers themselves in package inserts. **Since Roe v Wade in 1973 there have been over 50 million surgical abortions. It is now estimated that more than 250 million chemical abortions have occurred as result of women taking hormonal contraceptives.**⁵
- Today, for every child born in the United States, there are an estimated two children killed in the womb from surgical abortion and abortifacient contraceptives.⁶
- **98% of all women in the U.S. who have had sexual intercourse have used at least one contraceptive method.**⁷
- Since the advent of 'The Pill' in 1960, the average number of children in the American Catholic family has fallen from 5.5 to 2.1.⁸
- By age 45, at least one partner in every two U.S. marriages has been sterilized.⁹



- Only one in three U.S. couples now reaches their 25th wedding anniversary; only one in five reaches their 35th.¹⁰
- **The divorce rate for U.S. couples who use natural family planning to space children is less than 1%.**¹¹

SOURCES

1. *Sexually Transmitted Diseases Surveillance and Statistics*, U.S. Center for Disease Control and Prevention, 2004
2. Kahlenborn et al., "Oral Contraceptive Use - as a risk factor for premenopausal breast cancer: a meta analysis", Mayo Clinic proceedings 2006, pp. 1290 – 1320, 2006.
3. Bosch, et al., "Effect of oral contraceptives on risk of cervical cancer ...", International Agency of Research on Cancer, 2002
4. International Agency for Research on Cancer; research arm of the World Health Organization, 2010
5. Kuhar et al., "Infant Homicides through Contraceptives", International Pharmacists for Life, 5th edition, 2003
6. National Center for Health Statistics Division of Center for Disease Control, 2003. International Pharmacists for Life, 2003
7. Schneier, "Reaching the Cafeteria Catholic", 2006. The National Survey of Family Growth, U.S. Department of Health and Human Services, 2005
8. Kenneth C. Jones; "Index of Leading Catholic Indicators: the Church since Vatican II; *Catholic World Report*; May 2003
9. "Method of contraception use among women aged 18-44 currently using contraception, by marital status", Centers for Disease Control and Prevention, National Center for Health, 2002
10. U.S. Census Bureau, 2005
11. Dr. Robert Lerner, University of Chicago, "National Survey of Family Growth", U.S. Government's National Center for Health Statistics and "General Social Survey", National Opinion Research Center, 2000. Physicians for Life, 2007